

SÉMINAIRE DE THÉORIE SPECTRALE ET GÉOMÉTRIE

URSULA HAMENSTÄDT

Some aspects of the Laplace operator in negative curvature

Séminaire de Théorie spectrale et géométrie, tome S9 (1991), p. 95-97

http://www.numdam.org/item?id=TSG_1991__S9__95_0

© Séminaire de Théorie spectrale et géométrie (Grenoble), 1991, tous droits réservés.

L'accès aux archives de la revue « Séminaire de Théorie spectrale et géométrie » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

**RENCONTRES DE THEORIE SPECTRALE ET GEOMETRIE
GRENOBLE 1991
(Aussois du 7 au 14 avril)**

**Some aspects of the Laplace operator
in negative curvature**

Ursula HAMENSTÄDT

**Mathematisches Institut
der Universität Bonn
Beringstrasse 1
D-W 5300 BONN 1
ALLEMAGNE**

Let M be a compact Riemannian manifold of negative sectional curvature with universal covering \widetilde{M} . This covering is diffeomorphic to \mathbf{R}^n and admits a natural compactification by adding a sphere $\partial\widetilde{M}$ at infinity. The Wiener measure on paths induces for every $x \in \widetilde{M}$ a probability measure ω^x on $\partial\widetilde{M}$ which can naturally be considered as a Borel-probability measure on the fibre $T_x^1\widetilde{M}$ at x of the unit tangent bundle $T^1\widetilde{M}$ of \widetilde{M} ([P], [A], [AS]). These measures are equivariant under the action of the fundamental group Γ of M on $T^1\widetilde{M}$ and hence we can define a Borel-probability measure ω^s on the unit tangent bundle T^1M of M by $\omega^s(A) = \int_{x \in M} \omega^x(A \cap T_x^1M) dx$ (here dx is the normalized Lebesgue measure).

Recall that T^1M admits foliations W^i which are invariant under the action of the geodesic flow Φ^t ($i = s, ss$), the *stable foliation* W^s and the *strong stable foliation* W^{ss} . Each leaf of W^s is locally diffeomorphic to M and hence the Riemannian metric on M lifts to a Riemannian g^s on the leaves of W^s . The Laplace operator on the leaves with respect to this metric then induces a globally defined second order differential operator Δ^s on T^1M with continuous coefficients.

LEDRAPPIER. — *The measure ω^s is the unique harmonic measure for Δ^s , i.e. the unique measure such that $\int \Delta^s \varphi d\omega^s = 0$ for all smooth functions φ on T^1M (up to a constant, see [L3]).*

Similarly we obtain an operator Δ^{ss} for W^{ss} and an invariant measure ω^{ss} . The projection of ω^{ss} is in the Lebesgue measure class and its conditionals on the fibres of $T^1M \rightarrow M$ are the (non-normalized) Patterson-Sullivan measures ([L3], [Kn], [Y]).

Let λ be the Lebesgue-Liouville measure on T^1M . Call M *asymptotically harmonic* if $\omega^{ss} = \lambda$.

Equivalent are (see [L1],[L2], [L3]) :

- i) M is asymptotically harmonic.
- ii) $\omega^s = \lambda$.
- iii) $\omega^s = \omega^{ss}$.
- iv) For every Busemann function θ on \widetilde{M} , the function $e^{-h\theta}$ is minimal harmonic.
- v) Let $G(x, y)$ be the Green's function on \widetilde{M} . Then there are positive constants $C > 0, h > 0$ such that

$$\lim_{d(x,y) \rightarrow \infty} G(x, y) e^{hd(x,y)} = C .$$

Here the constant h equals the topological entropy of the geodesic flow Φ^t on T^1M .

vi) The top of the L^2 -spectrum of Δ on \widetilde{M} equals $-\frac{h^2}{4}$. (This and related results can be found implicitly in [L1], [L3] and [H4]).

For $\dim M \leq 4$, locally symmetric spaces are the only asymptotically harmonic ones ([H4], [L4]). In fact in certain cases more can be said :

vii) If $\dim M = 2$ and if any two of the measures ω^s, ω^{ss} and λ are equivalent (i.e. if they have the same measure zero sets) then M is asymptotically harmonic (and hence has constant curvature) ([K1], [K2], [L2], [H3]).

viii) If ω^{ss} and λ are equivalent then M is asymptotically harmonic ([H4]).

CONJECTURE. — If any two of the measures $\omega^s, \omega^{ss}, \lambda$ are equivalent then M is locally symmetric.

References

- [A] ANCONA A. — *Negatively curved manifolds, elliptic operators and Martin boundary*, Ann. of Math., 125 (1987), 495–536.
- [AS] ANDERSON M., SCHOEN R. — *Positive harmonic functions on complete manifolds of negative curvature*, Ann. of Math., 121 (1985), 429–461.
- [H1] HAMENSTÄDT U. — *A new description of the Bowen-Margulis measure*, Erg. Th. Dynam. Sys., 9 (1989), .
- [H2] HAMENSTÄDT U. — *An explicite description of harmonic measure*, Math. Zeit., 205 (1990), 287–299.
- [H3] HAMENSTÄDT U. — *Time preserving conjugacies of geodesic flow*, to appear in Erg. Th. Dynam. Sys..
- [H4] HAMENSTÄDT U. — in preparation..
- [Kai] KAĪMANOVICH V. A. — *Invariant measures of the geodesic flow and measures at infinity on negatively curved manifolds*, Ann. IHP Physique Théorique, 1990.
- [K1] KATOK A. — *Entropy and closed geodesics*, Erg. Th. Dynam. Sys., 2 (1982), 339–365.
- [K2] KATOK A. — *Four applications of conformal equivalence to geometry and dynamics*, Erg. Th. Dynam. Sys., 8 (1988), 139–152.
- [Kn] KNIEPER G. — *Horospherical measure and rigidity of manifolds of negative curvature*, Preprint, 1991.
- [L1] LEDRAPPIER F. — *Ergodic properties of Brownian motion on covers of compact negatively curved manifolds*, Bol. Soc. Bras. Mat., 19 (1988), 115–140.
- [L2] LEDRAPPIER F. — *Harmonic measures and Bowen-Margulis measures*, Israel Journal of Math., 71 (1990), 275–287.
- [L3] LEDRAPPIER F. — *Ergodic properties of the stable foliation*, Preprint.
- [M1] MARGULIS G.A. — *Applications of ergodic theory to the investigation of manifolds of negative curvature*, Funct. Anal. Appl., 3 (1969), 335–336.
- [M2] MARGULIS G.A. — *Certain measures associated with U-flows on compact manifolds*, Funct. Anal. Appl., 4 (1970), 55–67.
- [P] PRAT J.J. — *Etude asymptotique et convergence angulaire du mouvement brownien sur une variété à courbure négative*, CRAS Paris, 280 (1985), A1539–A1542.
- [S] SULLIVAN D. — *Related aspects of positivity in Riemannian geometry*, J. Differential Geom., 25 (1987), 327–51.
- [Y] YUE C.B. — *Brownian motion on Anosov foliations, integral formula and rigidity*, Preprint.