

SÉMINAIRE DE THÉORIE SPECTRALE ET GÉOMÉTRIE

ALAIN VALETTE

An application of Ramanujan graphs to C^* -algebra tensor products, II

Séminaire de Théorie spectrale et géométrie, tome 14 (1995-1996), p. 105-107

http://www.numdam.org/item?id=TSG_1995-1996__14__105_0

© Séminaire de Théorie spectrale et géométrie (Grenoble), 1995-1996, tous droits réservés.

L'accès aux archives de la revue « Séminaire de Théorie spectrale et géométrie » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

AN APPLICATION OF RAMANUJAN GRAPHS TO C*-ALGEBRA TENSOR PRODUCTS, II

Alain VALETTE

Let A, B be C*-algebras; denote by $A \otimes B$ the algebraic tensor product of A and B . According to the general theory of tensor products of C*-algebras, C*-norms on $A \otimes B$ lie between a minimal C*-norm, denoted by $\|\cdot\|_{\min}$, and a maximal C*-norm, denoted by $\|\cdot\|_{\max}$ (see [Tak79] for all this). Denote by H an infinite-dimensional separable Hilbert space, and by $B(H)$ the C*-algebra of linear, bounded operators on H . It was an old problem to decide whether or not all C*-norms coincide on $B(H) \otimes B(H)$; in a remarkable paper [JP95], M. Junge et G. Pisier give a negative answer to that question. More precisely, to describe quantitatively the discrepancy between the maximal and minimal C*-norms on $B(H) \otimes B(H)$, they introduce for any $n \in \mathbb{N}$ the number

$$\lambda(n) = \sup \left\{ \frac{\|u\|_{\max}}{\|u\|_{\min}} : u \text{ tensor of rank } n \text{ in } B(H) \otimes B(H) \right\}$$

and they prove:

THEOREM 1. — *There exists a constant $c > 0$ such that, for any $n \in \mathbb{N}$:*

$$c\sqrt[n]{n} \leq \lambda(n) \leq \sqrt{n}.$$

In [JP95], Junge and Pisier ask for the precise asymptotic behaviour of $\lambda(n)$ for $n \rightarrow \infty$. Actually they provide an excellent way of getting lower bounds on $\lambda(n)$ by introducing another constant C_n as follows:

DEFINITION 1. — *Let \mathbf{F}_n denote the free group on n generators a_1, a_2, \dots, a_n . Let C_n denote the infimum of all numbers $C > 0$ for which there exists a sequence $(\pi_k)_{k \geq 1}$ of unitary, finite-dimensional representations of \mathbf{F}_n such that:*

$$\left\| \sum_{i=1}^n (\pi_k \otimes \overline{\pi_m})(a_i) \right\| \leq C \text{ for all } k \neq m$$

(where $\overline{\pi_m}$ denotes the contragredient representation of π_m).

It is then proved in [JP95] that the constants $\lambda(n)$ and C_n are related through the following inequality:

$$\frac{n}{C_n} \leq \lambda(n) \quad (*)$$

In [Pis], Pisier proved:

PROPOSITION 1. — For any $n \geq 2$, one has

$$C_n \geq 2\sqrt{n-1}.$$

On the other hand, using Ramanujan graphs (see [Lub94]), I proved in [Val]:

PROPOSITION 2. — Let q be a prime power. Then

$$C_{q+1} \leq 2\sqrt{q}.$$

From these two results, it is natural to conjecture that $C_n = 2\sqrt{n-1}$ for any $n \geq 2$. I shall confirm this conjecture by proving that it holds asymptotically. I am grateful to U. Haagerup and G. Skandalis for suggesting to me the possibility of such a proof.

PROPOSITION 3.

$$\lim_{n \rightarrow \infty} \frac{C_n}{2\sqrt{n}} = 1$$

Proof: It follows from Proposition 3 in [Val] that the sequence $(\frac{C_n}{2\sqrt{n}})_{n \geq 1}$ is bounded, so let K be an upper bound for that sequence. It is also known that the function $n \rightarrow C_n$ is sub-additive (see the lemma in [Val]). Denote by p_k the k -th prime. For fixed n , let k be such that $p_k + 1 \leq n < p_{k+1}$. Then:

$$C_n \leq C_{p_k+1} + C_{n-p_k-1}$$

Using Proposition 2, we get:

$$C_n \leq 2\sqrt{p_k} + 2K\sqrt{n-p_k-1} \leq 2\sqrt{p_k} + 2K\sqrt{p_{k+1}-p_k}$$

Hence:

$$\frac{C_n}{2\sqrt{n}} \leq \frac{C_n}{2\sqrt{p_k}} \leq 1 + K\sqrt{\frac{p_{k+1}}{p_k} - 1}.$$

By the prime number theorem of Hadamard and de la Vallée-Poussin, the ratio $\frac{p_{k+1}}{p_k}$ tends to 1 for $k \rightarrow \infty$. Thus:

$$\limsup_{n \rightarrow \infty} \frac{C_n}{2\sqrt{n}} \leq 1.$$

On the other hand, it follows from Proposition 1 that we also have

$$\liminf_{n \rightarrow \infty} \frac{C_n}{2\sqrt{n}} \geq 1,$$

so that the result is proved.

From Proposition 3 and the inequality (*), it immediately follows that:

COROLLARY 1. —

$$\liminf_{n \rightarrow \infty} \frac{\lambda(n)}{\sqrt{n}} \geq \frac{1}{2}.$$

References

- [JP95] M. Junge and G. Pisier. Bilinear forms on exact operator spaces and $B(H) \otimes B(H)$. *Geometric and Functional Analysis*, 5:329–363, 1995.
- [Lub94] A. Lubotzky. *Discrete groups, expanding graphs and invariant measures*. Progress in Math. 125, Birkhäuser, 1994.
- [Pis] G. Pisier. Quadratic forms in unitary operators. Preprint, 1995.
- [Tak79] M. Takesaki. *Theory of operator algebras I*. Springer-Verlag, 1979.
- [Val] A. Valette. An application of Ramanujan graphs to C^* -algebra tensor products. To appear in *Discrete Math*.